Levine, J. F., 931
Levy, M. G., 931
Linley, J. R., 9, 578
Linley, P. A., 578
Linthicum, K. J., 537
Liscia, A., 14
Litwin, T. S., 945
Logan, T. M., 537
Longnecker, M. T., 632
Lopez-W., R., 440
Lorenz, L. H., 856, 878
Lounibos, L. P., 578
Lundstrom, J. O., 266, 561

M
Ma, M., 564
MacLean, D. C., 945
Main, A. J., 763, 968, 986
Maina, J. N., 316
Maire, A., 835
Mangan, R. L., 295
Mansour, M. M., 968, 982, 986
Manweiler, S. A., 1011
Marchiondo, A. A., 913
Marquino, M., 202
Marzouk, A. S., 288
Marzouk, A. S., 763
Mather, M. E., 646
Mather, T. N., 646
Matuschka, F.-R., 385
McHugh, C. P., 1027, 1065
McKinnon, C. N., 934
McLaughlin, R. E., 1031
McMeans, M. C., 1065
Mecham, J. O., 602
Meola, R. W., 1045
Messersmith, M., 1071
Meyer, R. P., 607
Meyer, R. P., 356
Milby, M. M., 811
Miller, J., 72
Miller, L. H., 905
Miyagi, I., 344, 530
Mogi, M., 509, 530
Mohamed, F. S. A., 288
Moon, R. D., 391, 727
Morales, A., 1
Morales-Malacara, J. B., 440
Morris, T. A., 543
Morrison, M. L., 1011
Moulton, J. R., 537
Mount, G. A., 543, 671, 686
Mulla, M. S., 57, 429
Mullins, D. E., 703
Mumcuoglu, K. Y., 72

N
Nagardeolekar, M., 51
Nasci, R. S., 716, 1031
Nayar, J. K., 409
Needham, G. R., 651
Nesbitt, S. A. T., 586
Nicholson, W. L., 931
Niklasson, B., 266, 561
Noda, N., 501
Nokoe, S., 316
Nunamaker, R. A., 602

O
Oelofsen, M. J., 939
Ohmori, D., 302
Ohtomo, H., 501
Okazawa, T., 446
Oliver, Jr., J. H., 324
Omar, B., 851
Omar, S., 851
Onyango, F. K., 377, 570
Osburn, B. I., 331
Oster, C. N., 570

P
Palau, M. T., 1
Palma, K. G., 1045
Pandey, K. S., 962
Pasteur, N., 202
Patel, K. J., 892
Patterson, R. S., 1062
Perkins, P. V., 377, 570
Petrarca, V., 564
Peyton, E. L., 207
Pfuntner, A. R., 811
Pickel, S. L., 632
Pietra, P., 14
Pilakasiri, C., 1016
Powell, D. G., 874
Prakash, S., 962
Presley, S. M., 750
Presser, S. B., 811, 1008
Punyua, D. K., 313

R
Rae, D. J., 756
Rao, K. M., 962
Rapp, C. M., 1035
Rashed, S. S., 429
Ratzlaff, R. E., 1002
Reifenrath, W. G., 494
Reisen, W. K., 356, 607, 811
Richter, D., 385
Riner, J. L., 913
Roberts, A. W., 874
Roberts, C. R., 681
Roe, R. M., 36
Rolseth, B. M., 586
Romans, P., 905
Rosen, L. J., 72
Roshdy, M. A., 288, 763
Ross, M. H., 703
Rowe, K. F., 913
Rueda, L. M., 892
Runstrom, E. S., 515
Rutledge, L. C., 406, 494
Rutz, D. A., 309

S
Sabattini, M. S., 784
Sadler, J. P., 628
Said, S. M. E., 773
Sakai, R. K., 905
Samuel, W. M., 656
Sarojo, U. T. I. B., 777
Sattabongkot, J., 1016
Sawyer, M. M., 331
Scholl, P. J., 523
Schriefer, M. E., 36
Schwan, T. G., 720
Scott, T. W., 856, 864, 878, 999
Seabury, F., 9
Sereno, R., 784
Shanbaky, N. M., 968, 975, 982, 986
Shehata, M. G., 592, 773
Sherif, N. E.-D. H. S., 651
Shoumar, N. F., 592
Shroyer, D. A., 334
Siegel, J., 556
Slusser, J. H., 913
Smith, M. S., 939
Snoddy, E. L., 543
Sonenshine, D. E., 36, 763, 913
Song, Q., 564
Spielman, A., 196, 385, 945
Spoehel, J. J., 356
Srivastava, C. P., 962
Steinkraus, D. C., 309
Stinner, R. E., 892, 922
Stoffolano, Jr., J. G., 14
Stricker, K. W., 51
Stubbs, H. A., 282
Sulaiman, S., 851
Suleman, M., 819
Suwanpanit, P., 509
Suzuki, K., 302
Szumlas, D. E., 480

T
Tabachnick, W. J., 24
Takaoka, H., 701
Tanaka, K., 344
Teel, P. D., 632
Tempelis, C. H., 356, 811
Thomas, R. E., 720
Tietze, N. S., 57
Todd, Jr., K. S., 556
Todd, L., 955
Toma, T., 344
Tovey, E. R., 368
Trenggono, B., 777
Tsuji, N., 446
Turell, M. J., 266
Tyler, R. D., 803

U
Uchida, K., 302
Uehara, K., 993

V
Van der Linde, T. C. de K., 939
Van Riper III, C., 68
Venkatesh, K., 36
Vickery, A. C., 409

W
Wagner, G. G., 1067
Walker, A. R., 955
Walton, T. E., 487
Walton, W. E., 57
Washino, R. K., 68, 331
Watts, D. M., 537
Weatherhead, A., 1016
Weaver, S. C., 864, 878
Weintraub, J., 523
Welch, D. A., 656

Welch, J. B., 295
Wellems, T. E., 905
Whitmire, R. E., 570
Wigington, J. G., 602
Wikel, S. K., 1002
Wilkerson, R. C., 207
Williams, N. M., 874
Wilson, M. L., 945
Windsor, I. M., 697
Wirth, M. C., 202
Wirtz, R. A., 377, 1016
Wooster, M. T., 703
Work, T. M., 68, 331
Wright, R. E., 51, 480

Y
Yamakura, F., 302
Yamamura, N., 446
Young, D. G., 1
Yunker, C. E., 651
Yuval, B., 196

Z
Zhang, J.-Z., 564
Zimmerman, R. H., 543

CUMULATIVE SUBJECT INDEX
Journal of Economic Entomology
Volume 83, 1990

A
abamectin, bioassays, 1935
Blattella germanica, 1243
cross-resistance, 1243
insecticide selectivity, 1783
integrated pest management, 1783
Lymantria dispar, 917
MK-243, 917
Musca domestica, 1935
acaricides, *Panonychus*, 1711
Tetranychus, 1711
acceptance, *Empoasca fabae*, 814
settling, 814
Acrosternum hilare, cotton, 842
yield reduction, 842
action threshold, aphids, 1381
integrated pest management, 1381
activity, actograph, 2385
Conotrachelus nenuphar, 2385
actograph, activity, 2385
Conotrachelus nenuphar, 2385
adhesives, Stomoxys calcitrans, 883
traps, 883
Adoretus sinicus, carbohydrates, 2022
feeding preference, 2022
aerial deposition, *Bacillus thuringiensis*, 173
gypsy moths, 173
aesthetic injury levels, *Anisota senatoria*, 2044
Cecidomyia spp., 485
dispersion, 485
insect control, 2044
age distribution, *Cochliomyia hominivorax*, 1422
pteridine eye pigments, 1422
aggregation pheromones, insect traps, 273
Oryzaephilus surinamensis, 273
Agrotis ipsilon, corn seedling damage, 1058
laboratory bioassays, 2073
Pseudaletia unipuncta, 596
pyrethroids, 2073
sampling, 596
weed competition, 1058
alfalfa, cutting, 2427
economic injury levels, 1587, 1612
Empoasca fabae, 807, 1587
Otiorhynchus ligustici, 1612
translocation disruption, 807
alfalfa hay, alfalfa insects, 2050
clear-cutting, 2050
alfalfa insects, alfalfa hay, 2050
clear-cutting, 2050
alginate, *Beauveria bassiana*, 2225
cereal aphids, 2225
almonds, *Amyelois transitella*, 570
population monitoring, 570
amino acid, Ninhydrin, 1633
spray deposition, 1633
Amyelois transitella, almonds, 570
population monitoring, 570
Anarsia lineatella, distribution, 505
sampling, 505
Anasa tristis, Cucurbitaceae, 943
insect-plant interactions, 943
mulch, 1988
vegetable production systems, 1988
Anastrepha, *Ceratitis*, 1940
quarantine treatments, 1940
Anastrepha ludens, color, 412
trapping, 412
Anastrepha suspensa, carambolas, 458
commodity treatment, 458, 1471, 1475, 2340
fruit quality, 1468
heat treatment, 2331
Mangifera indica, 2320
quarantine security, 1468, 1471, 1475, 2320, 2331, 2340
quarantine treatments, 2324
shrink wrapping, 2324
Ancylis comptana, oviposition, 1514
strawberry, 1514
Anisopteromalus calandrae, biological control, 2462
Sitophilus zeamais, 2462
Anisota senatoria, aesthetic injury level, 2044
insect control, 2044
Anobiidae, Dermestidae, 879
museum pest control, 879
Anophothrips obscurus, *Frankliniella fusca*, 621
Frankliniella tenuicornis, 621
Anoplura, Mallophaga, 1435
sampling, 1435
Anthonomus eugenii, *Capsicum annuum*, 2003
pest management, 2003
Anthonomus grandis grandis, egg hatch, 1853
emergence, 1879
pheromone, 610
sterility, 1853
survival, 1879
Anthonomus grandis, predation, 1
Solenopsis invicta, 1
Anthonomus quadrigibbus, *Conotrachelus crataegi*, 1326
Conotrachelus nenuphar, 1326
anthurium, floriculture, 1531
integrated pest management, 1531
antibiosis, *Heliothis zea*, 614
plant resistance, 1053
population reduction, 614
Sipha flava, 1053
Anticarsia gemmatalis, *Glycine max*, 2107
host plant resistance, 2107
antidesiccants, *Heterorhabditis*, 1795
Steinernema, 1795
Aonidiella aurantii, insecticides, 189
phenology, 189
Aphidoletes aphidimyza, cold storage, 965
diapause, 965
aphids, action threshold, 1381
asparagus, 1994
Brachycorynella asparagi, 1994
filberts, 41
insecticde resistance, 41
integrated pest management, 1381
Aphis gossypii, bifenthrin, 955
videotape recording, 955
Apis mellifera, citrus, 1267
colony strength, 748
detoxification, 715
digitizer, 1791
dimethoate, 1267
feral honey bee colonies, 81
Galleria mellonella, 760, 766
honey bees, 81
knockdown, 1925
microbial control, 760, 766
microbial safety test, 755
nontarget organism, 755
Orthoptera, 784
packaged bees, 1271
pheromones, 1271
population dynamics, 1791
population estimates, 748
Poxviridae, 784
resistance, 715
wasp and hornet sprays, 1925
apple pests, integrated pest management, 2405
Rhagoletis pomonella, 2405
apple volatiles, *Rhagoletis pomonella*, 539
trapping, 539
apples, control, 2401
Cydia pomonella, 466
economic injury level, 552
Magicicada septendecim, 2401
Panonychus ulmi, 552
Prunus, 466

Arachnida, bioassays, 8
pesticides, 8
asparagus, aphids, 1994
Brachycorynella asparagi, 1994
attractancy, color preference, 111
Macrodactylus subspinosus, 111
attractants, *Ceratitis capitata*, 819, 1350
Cetoniinae, 1298
curcurbitacin, 1085
Diabrotica virgifera virgifera, 1085
lure, 1350
Rutelinae, 1298
trimedlure, 819
attraction, semiochemicals, 2040
Toxotrypana curvicauda, 2040
avermectin, fenbutatin oxide, 1752
hexythiazox, 1752
Lymantria dispar, 710
sublethal, 710
azadirachtin, metamorphosis, 2168
Tephritidae, 2168

B
Bacillus sphaericus, *Bacillus thuringiensis israelensis*, 347, 1280
Culicidae, 1280
Mammalia, 347
Bacillus thuringiensis, aerial deposition, 173
formulation, 1813
gypsy moth maternal effects, 2211
gypsy moths, 173
insecticide resistance, 1671
microbial pesticide, 1671
microbial pesticide resistance, 2211
Operophtera brumata, 920
Ostrinia nubilalis, 1813
Panonychus ulmi, 920
Tetranychus urticae, 792
thuringiensin, 792
Bacillus thuringiensis ssp *israelensis*, *Bacillus sphaericus*, 347 Mammalia, 347
Bacillus thuringiensis israelensis, *Bacillus sphaericus*, 1280 Culicidae, 1280
backcross sterility, genetic control, 405
tobacco budworm, 405
baculovirus, dyes, 168
ultraviolet, 168
baits, Diptera pupae, 67
Solenopsis invicta, 67
barrier treatments, *Iridomyrmex humilis*, 1402
repellency, 1402
Beauveria bassiana, alginate, 2225
cereal aphids, 2225
Chalcodermus bimaculatus, 1276
microbial control, 1276
behavior, insecticides, 217
pyrethroids, 217, 1510
Trichoplusia ni, 1510
behavioral resistance, insecticides, 1216
pyrethroids, 1216
Bemisia tabaci, buprofezin, 1254
esterase activity, 1665
insecticide resistance, 1665
vapor toxicity, 1254
benzodioxole, fruit flies, 1261
soil, 1261
Bethylidae, *Goniozus legneri*, 733
insecticidal toxins, 733
bifenthrin, *Aphis gossypii*, 955
videotape recording, 955
binomial sampling, *Panonychus ulmi*, 420
Tetranychus urticae, 420
bioassay, chemigation, 1049
Ostrinia nubilalis, 1049
bioassay methods, *Lygus* spp., 2200
thuringiensin, 2200
bioassays, abamectin, 1935
Arachnida, 8
Musca domestica, 1935
pesticides, 8
biological control, *Anisopteromalus calandrae*, 2462
Cylas formicarius elegantulus, 1818
diapause, 2196
entomogenous nematodes, 97, 2380
Exeristes roborator, 126
learning, 126
Liriomyza trifolii, 2380
Lymantria dispar, 2229
Mayetiola destructor, 2269
mirid pests, 101
Nematoda, 1286
nematodes, 1818
parasites, 101, 1096
parasitoids, 2269
predators, 2229
Pyrrhalta, 2196
Rhabditida, 97
Sitophilus zeamais, 2462
Steinernema feltiae, 1286
stored-product insects, 1096
Biosteres tryoni, *Ceratitis capitata*, 779
sex ratio, 779
biotic potential, food preference, 1338
Melanoplus sanguinipes, 1338
biotypes, *Daktulosphaira vitifoliae*, 489
grape phylloxera, 489
bipyridylium herbicides, 20–OH ecdysone, 2175
Neobellieria bullata, 2175
birdsfoot trefoil, insect control, 2086
plant bugs, 580, 2086
plant damage, 580
Blatta orientalis, genitalia, 1415
hydroprene, 1415
Blattella germanica, abamectin, 1243
chlorpyrifos, 441, 1912
crack and crevice, 1907
cross-resistance, 1243
delayed-action toxicity, 148
dispersal, 142
Ebeling choice box, 153
fenoxycarb, 444
growth, 2295
insect growth regulators, 2295
insecticidal baits, 153
insecticides, 142, 2290
population effects, 444
propoxur, 1912
reproduction, 441
residual insecticides, 1907
sulfluramid, 148, 1409
toxicity, 1409, 2290
Blissus leucopterus, thatch, 2370, 2375
turfgrass, 2370, 2375
Blissus leucopterus hirtus, endophyte, 1640
plant resistance, 1640
Blissus leucopterus leucopterus, *Zea mays*, 618
blister beetle, cantharidin, 1042
Epicauta, 1042
blossom end defects, *Dacus*, 1883
infestation rates, 1883
boll weevils, reproduction, 392
Sphaeralcea spp., 392
bovine dung, Muscidae, 1446
nematodes, 1446
Brachycorynella asparagi, aphids, 1994
asparagus, 1994
Bracon hebetor, *Cadra cautella*, 1110
insect-resistant packaging, 1110
Brassica napus L., rapeseed, 2246
buprofezin, *Bemisia tabaci*, 1254
vapor toxicity, 1254

C
cabbage, *Plutella xylostella*, 1201
resistance, 1201

Cacopsylla pyricola, juvenile hormone, 293
overwintering, 293
pyrethroids, 2159
resistance, 2159
Cadra cautella, *Bracon hebetor*, 1110
insect-resistant packaging, 1110
Calocoris norvegicus, pistachios, 1841
weeds, 1841
Campylomma verbasci, pest management, 1506
pheromones, 1508
cantharidin, blister beetle, 1042
Epicauta, 1042
Capsicum annuum, *Anthonomus eugenii*, 2003
pest management, 2003
carambolas, *Anastrepha suspensa*, 458
commodity treatment, 458
carbamate, pyrethroids, 2136
Spodoptera exigua, 2136
carbaryl consumption, mortality, 2164
wild rodents, 2164
carbendazim, metabolism, 742
Rhizoglyphus echinopus, 742
carbohydrates, *Adoretus sinicus*, 2022
feeding preference, 2022
carbon dioxide, *Cryptolestes ferrugineus*, 277
wheat, 277
cattle, *Haematobia irritans*, 1430
integrated pest management, 1430
cauliflower, defoliation, 1499
Lepidoptera, 1499
Cecidomyia spp., aesthetic injury levels, 485
dispersion, 485
cell culture, cotton, 737
Heliothis virescens, 737
Ceratitis, host plant, 1863
infestation rate, 1863
quarantine treatments, 1940
Ceratitis, *Anastrepha*, 1940
Ceratitis capitata, attractant, 819, 1350
Biosteres tryoni, 779
diet, 1954
distribution, 1949
gamma irradiation, 1449
linalool, 2235
lure, 1350
protein, 1954
pyrazine, 2235
quarantine, 1449
sex ratio, 779
sperm, 1949
trimedlure, 819
cereal aphids, alginate, 2225
Beauveria bassiana, 2225
Diuraphis noxia, 846
host plants, 846
Cetoniinae, attractants, 1298
Rutelinae, 1298
Chalcodermus bimaculatus, *Beauveria bassiana*, 1276
microbial control, 1276
chemical adsorption, field burning, 55
pest control, 55
chemical control, corn, 1621
Keiferia lycopersicella, 212
Oligonychus pratensis, 1621
tomatoes, 212
chemical termination treatment, cotton, 1074
Pectinophora gossypiella, 1074
chemigation, bioassay, 1049
Ostrinia nubilalis, 1049
chlordimeform, cotton phytotoxicity, 1549
methyl parathion, 1549
chlorpyrifos, *Blattella germanica*, 441, 1912
Oligonychus pratensis, 1570
parathion, 1570
propoxur, 1912
reproduction, 441
chlorpyrifos methyl, malathion, 1636
rice, 1636
chlorpyrifos-methyl, granaries, 1128
oats, 1128
Choristoneura fumiferana, decision support system, 1479
Wald's seq. prob. ratio test, 1479
Choristoneura rosaceana, *Cydia pomonella*, 434
monitoring, 434
chrysanthemums, *Liriomyza trifolii*, 1246
neem seed extract, 1246
citrus, *Apis mellifera*, 1267
dimethoate, 1267
oviposition, 860
Pantomorus cervinus, 860
Citrus, gamma radiation, 905
Panonychus citri, 976
Pantomorus cervinus, 905
yield, 976
Clavigralla tomentosicollis, resistance, 300
wild cowpea, 300
clear-cutting, alfalfa hay, 2050
alfalfa insects, 2050
Coccinella septempunctata, *Hippodamia variegata*, 1292
Propylea quatuordecimpunctata, 1292
Cochliomyia hominivorax, age distribution, 1422
pteridine eye pigments, 1422
Cochylis hospes, Lepidoptera, 135
sunflowers, 135
cockroaches, resistance, 1698
synergist, 1698
codling moths, *Cydia pomonella*, 131
oviposition, 131
cold storage, *Aphidoletes aphidimyza*, 965
diapause, 965
Coleomegilla maculata, *Leptinotarsa decemlineata*, 1306
predation, 1306
Coleoptera, distribution, 1102
pheromones, 268
probe traps, 1102
stored grain, 268
Colladonus montanus, *Fieberiella florii*, 2279
western X-disease, 2279
colony strength, *Apis mellifera*, 748
population estimates, 748
color, *Anastrepha ludens*, 412
trapping, 412
color preference, attractancy, 111
Macrodactylus subspinosus, 111
color traps, cucumber, 971
Frankliniella occidentalis, 971
Colorado potato beetles, crop rotation, 315
insecticide resistance, 315
combined injuries, *Glycine max*, 599
Plathypena scabra, 599
commodity, *Quadraspidiotus perniciosus*, 892
radiation, 892
commodity treatment, *Anastrepha suspensa*, 1475
Anastrepha suspensa, 458, 1471, 2340
carambolas, 458
quarantine security, 1471, 1475, 2340
cone and seed insects, *Pseudotsuga menziesii*, 1485
western conifer seed bug, 1485
Conophthorus coniperda, seed orchards, 2349
white pine, 2349
Conotrachelus crataegi, *Anthonomus quadrigibbus*, 1326
Conotrachelus nenuphar, 1326
Conotrachelus nenuphar, activity, 2385
actograph, 2385
Anthonomus quadrigibbus, 1326
Conotrachelus crataegi, 1326

conservation tillage, *Ostrinia nubilalis*, 2455
continuous measure, *Heliothis virescens*, 1222
insecticide resistance, 1222
control, apples, 2401
Magicicada septendecim, 2401
Musca domestica, 788
Muscovy ducks, 788
controlled atmosphere, *Cydia pomonella*, 472
humidity, 472
controlled atmosphere storage, *Cylas formicarius elegantulus*, 461 *Ipomoea batatas*, 461
Coptotermes formosanus, repellency, 875
Reticulitermes flavipes, 1918
termiticides, 875
tunneling response, 1918
corn, chemical control, 1621
DIMBOA, 1626
economic injury, 2452
injury level, 1069, 1595
feeding behavior, 850
insecticides, 1114
Oligonychus pratensis, 1069, 1621
Ostrinia nubilalis, 1595
Rhopalosiphum maidis, 1626
orn, *Spodoptera frugiperda*, 2452
stalkborers, 850
stored-product insects, 1114
corn rootworms, *Diabrotica*, 574
insecticides, 226
residual, 226
sampling, 574
corn seedling damage, *Agrotis ipsilon*, 1058
weed competition, 1058
cotton, *Acrosternum hilare*, 842
cell culture, 737
chemical termination treatment, 1074
diapause, 1827
Frankliniella spp., 1064
Helicoverpa spp., 1827
Heliothis virescens, 737
Heliothis zea, 2115
Pectinophora gossypiella, 1074
sampling procedures, 1064
toxicity, 2115
yield reduction, 842
cotton phytotoxicity, chlordimeform, 1549
methyl parathion, 1549
crack and crevice, *Blattella germanica*, 1907
residual insecticides, 1907
critical radius, metabolic heat, 896
thermal conductivity, 896
crop rotation, Colorado potato beetles, 315
insecticide resistance, 315
cross-resistance, abamectin, 1243
Blattella germanica, 1243
fluvalinate resistance, 698
Scirtothrips citri, 698
Cryptolestes ferrugineus, carbon dioxide, 277
wheat, 277
Cryptostigmata, Kentucky bluegrass, 205
Poa pratensis, 205
Ctenocephalides felis felis, *Felis catus*, 2306
repellency, 2306
cucumbers, color traps, 971
damage, 2092
Frankliniella occidentalis, 971
thrips, 2092
Cucurbitaceae, *Anasa tristis*, 943
insect-plant interactions, 943
Culex quinquefasciatus, esterases, 1704
insecticide resistance, 1704
Culex spp., esterases, 2187
organophosphate resistance, 2187
Culicidae, *Bacillus sphaericus*, 1280
Bacillus thuringiensis israelensis, 1280
cultivar blends, cultivar mixtures, 255
cultural control, *Elasmopalpus lignosellus*, 1034
Labidura riparia, 1034
Mayetiola destructor, 1025
Triticum aestivum, 1025
curcurbitacin, attractants, 1085
Diabrotica virgifera virgifera, 1085
cutting, alfalfa, 2427
swathing, 2427
Cydia pomonella, apples, 466
Choristoneura rosaceana, 434
codling moths, 131
controlled atmosphere, 472
humidity, 472
monitoring, 434
oviposition, 131
postharvest quality, 2335
Prunus, 466
Prunus spp., 2335
Cylas formicarius elegantulus, biological control, 1818
controlled atmosphere storage, 461
Ipomoea batatas, 461, 1901
nematodes, 1818
sampling, 1901
Cynodon dactylon, resistance, 585
Spodoptera frugiperda, 585
cyromazine, *Delia antiqua*, 2398
diflubenzuron, 1689
growth regulator, 2398
resistance, 1689

D

Dacus, blossom end defects, 1883
infestation rates, 476, 1883
ripeness index, 476
Dacus cucurbitae, *Dacus dorsalis*, 2274
sampling, 2274
Dacus dorsalis, *Dacus cucurbitae*, 2274
heat, 160
papayas, 160, 1944
sampling, 2274
temperature stress, 1944
Dacus tryoni, kiwifruit, 2317
quarantine treatments, 2317
Dacus spp., papayas, 2327
quarantine treatment, 2327
Daktulosphaira vitifoliae, biotypes, 489
grape phylloxera, 494, 489
indoleacetic acid, 494
damage, cucumbers, 2092
degree-days, 837
thrips, 2092
Xanthogaleruca luteola, 837
date palms, *Ectomyelois ceratoniae*, 2357
nitidulid beetles, 2357
decision support system, *Choristoneura fumiferana*, 1479
Wald's seq.-prob. ratio test, 1479
defoliation, cauliflower, 1499
Lepidoptera, 1499
degree-days, damage, 837
Mayetiola destructor, 1015
phenology, 1015
sugarbeets, 2078
Tetanops myopaeformis, 2078
Xanthogaleruca luteola, 837
delayed-action toxicity, *Blattella germanica*, 148
sulfluramid, 148
Delia antiqua, cyromazine, 2398
insect growth regulator, 2398
Delia coarctata, expert system, 2065
pest management, 2065
Dermestidae, Anobiidae, 879
museum pest control, 879
detector dog, quarantine, 1932
screwworms, 1932
detoxification, *Apis mellifera*, 715
resistance, 715
detoxifying enzymes, feeding habits, 361
Diabrotica, corn rootworms, 574
sampling, 574

Diabrotica barberi, disubstituted benzenes, 1316
lure, 1316
Diabrotica undecimpunctata howardi, lectins, 2480
Ostrinia nubilalis, 2480
Diabrotica vergifera vergifera, enhanced degradation, 1250
fonofos, 1250
attractants, 1085
curcurbitacin, 1085
mark–release–capture, 1360
tethered-flight assay, 1360
Diadegma terebrans, hydroxamic acids, 356
Ostrinia nubilalis, 356
diagnostic dose, resistance, 1151, 1160
statistical power, 1160
statistical test power, 1151
diapause, *Aphidoletes aphidimyza*, 965
biological control, 2196
cold storage, 965
cotton, 1827
Helicoverpa spp., 1827
Pyrrhalta, 2196
Diatraea grandiosella, oviposition, 1575
plant resistance, 1578
sampling, 1370
Spodoptera frugiperda, 1575, 1578
Taylor's power law, 1370
Diatraea saccharalis, host plant resistance, 221
integrated pest management, 1563
plant resistance, 866, 1563
pubescence, 866
sugarcane, 221
dieldrin, *Drosophila melanogaster*, 1733
insecticide resistance, 1733
diet, *Ceratitis capitata*, 1954
protein, 1954
diflubenzuron, cyromazine, 1689
resistance, 1689
digitizer, *Apis mellifera*, 1791
population dynamics, 1791
DIMBOA, corn, 1626
Rhopalosiphum maidis, 1626
dimethoate, *Apis mellifera*, 1267
citrus, 1267
Diptera pupae, baits, 67
Solenopsis invicta, 67
disinfestation, irradiation, 1435
Tetranychus urticae, 1455
disparlure, insect pheromone, 910
Lymantria dispar, 910, 1977
sex pheromones, 1977
dispersal, *Blattella germanica*, 142
insecticides, 142
Ostrinia nubilalis, 831
scouting, 831
dispersion, aesthetic injury levels, 485
Cecidomyia spp., 485
pest resurgence, 1847
Tetranychus urticae, 1847
disruption, *Orgyia pseudotsugata*, 1487
pheromones, 1487
distribution, *Anarsia lineatella*, 505
Ceratitis capitata, 1949
Coleoptera, 1102
fumigant, 1459
phosphine, 1459
probe traps, 1102
sampling, 505
sperm, 1949
disubstituted benzenes, *Diabrotica barberi*, 1316
lure, 1316
Diuraphis noxia, cereal aphids, 846
host plants, 846
resistance, 1091
Triticum aestivum, 2434
vernalization, 2434
wheat, 1091
dose mortality, extravariation, 1649
goodness-of-fit, 1649
Drosophila melanogaster, dieldrin, 1733
insecticide resistance, 1733
mutagenicity, 721
permethrin, 721
dyes, baculovirus, 168
ultraviolet, 168

E

ear tags, *Haematobia irritans*, 1715
pyrethroid resistance, 1715
Ebeling choice box, *Blattella germanica*, 153
insecticidal baits, 153
economic injury, corn, 2452
Spodoptera frugiperda, 2452
economic injury levels, apples, 552
alfalfa, 1587, 1612
corn, 1069, 1595
Empoasca fabae, 1587
Leptinotarsa decemlineata, 2058
limes, 2008
Oligonychus pratensis, 1069
Ostrinia nubilalis, 1595
Otiorhynchus ligustici, 1612
Panonychus ulmi, 552
Polyphagotarsonemus latus, 2008
potatoes, 2058
Sipha flava, 1008
sorghum, 1008
economic threshold, integrated pest management, 641
soybeans, 641
Ectomyelois ceratoniae, date palms, 2357
nitidulid beetles, 2357
egg hatch, *Anthonomus grandis grandis*, 1853
sterility, 1853
Elasmopalpus lignosellus, cultural control, 1034
Labidura riparia, 1034
elm, insecticide bark bands, 1495
Xanthogaleruca luteola, 1495
emergence, *Anthonomus grandis grandis*, 1879
survival, 1879
Empoasca fabae, acceptance, 814
alfalfa, 807, 1587
economic injury levels, 1587
immigration, 1858
low temperature, 1541
Medicago sativa, 1858
mortality response, 1541
settling, 814
translocation disruption, 807
endophyte, *Blissus leucopterus hirtus*, 1640
plant resistance, 1640
enhanced degradation, *Diabrotica vergifera vergifera*, 1250
fonofos, 1250
entomogenous nematodes, biological control, 97, 2380
Liriomyza trifolii, 2380
Rhabditida, 97
Eotetranychus williamettei, *Tetranychus pacificus*, 2252
Vitis vinifera, 2252
Epicauta, blister beetle, 1042
cantharidin, 1042
Epiphyas postvittana, host-insect interaction, 2124
resistance, 2124
eradication, population management, 1344
quarantine, 1344
Eriosoma lanigerum, insect-plant interactions, 1526
integrated pest management, 1526
esterase activity, *Bemisia tabaci*, 1665
insecticide resistance, 1665
esterases, *Culex quinquefasciatus*, 1704
Culex spp., 2187

insecticide resistance, 1704
Lygus hesperus, 725
Myzus nicotianae, 2130
organophosphate resistance, 2187
resistance, 725, 2130
European red mite, *Panonychus ulmi*, 547
stress, 547
Exeristes roborator, biological control, 126
learning, 126
expert system, *Delia coarctata*, 2065
pest management, 2065
extravariation, dose mortality, 1649
goodness-of-fit, 1649

F
fecundity, herbivory, 377
longevity, 1333
necrophagy, 377
pallidovirens, 1333
feeding, grasshoppers, 2421
kochia, 2421
feeding behavior, corn, 850
Schizaphis graminum, 241
Sorghum bicolor, 241
stalkborers, 850
feeding deterrents, repellents, 629
Rhyzopertha dominica, 629
feeding duration, *Lygus lineolaris*, 2443
population density, 2443
feeding habits, detoxifying enzymes, 361
feeding preference, *Adoretus sinicus*, 2022
carbohydrates, 2022
feeding stimulant, *Ostrinia nubilalis*, 2207
starch encapsulation, 2207
Felis catus, *Ctenocephalides felis felis*, 2306
catus, repellency, 2306
fenbutatin oxide, avermectin, 1752
hexythiazox, 1752
fenoxycarb, *Blattella germanica*, 444
hydramethylnon, 74
megacephala, 74
population effects, 444
feral honey bee colonies, *Apis mellifera*, 81
honey bees, 81
Fieberiella florii, *Colladonus montanus*, 2279
western X-disease, 2279
field burning, chemical adsorption, 55
pest control, 55
field control, Pseudoplusia includens, 27
soybeans, 27
filberts, aphids, 41
insecticide resistance, 41
flies, poultry, 451
predators, 451
floriculture, anthurium, 1531
integrated pest management, 1531
flower thrips, peach injury, 511
sampling techniques, 511
fluvalinate resistance, cross-resistance, 698
Scirtothrips ciri, 698
fonofos, *Diabrotica vergifera vergifera*, 1250
enhanced degradation, 1250
food preference, biotic potential, 1338
Melanoplus sanguinipes, 1338
foraging, grasses, 260
Lotus corniculatus, 260
formamidine, house fly, 2181
synergism of pyrethroids, 2181
formulation, *Bacillus thuringiensis*, 1813
Ostrinia nubilalis, 1813
Frankliniella fusca, *Anophothrips obscurus*, 621
Frankliniella tenuicornis, 621
Frankliniella occidentalis, color traps, 971
cucumber, 971
integrated pest management, 1519
Thrips palmi, 1519
Frankliniella spp., cotton, 1064
sampling procedures, 1064
Frankliniella tenuicornis, *Anophothrips obscurus*, 621
Frankliniella fusca, 621
fruit flies, benzodioxole, 1261
heat, 1959
methoprene, 901
papaya, 901
quarantine treatments, 1959, 1965
soil, 1261
vapor heat, 1965
fruit orchards, pheromone traps, 197
trap efficiency, 197
fruit quality, *Anastrepha suspensa*, 1468
quarantine security, 1468
fumigants, distribution, 1459
phosphine, 1459
Rhyzopertha dominica, 1677
Tribolium castaneum, 1677
fungicide, *Nomuraea rileyi*, 2193
soybean defoliators, 2193
furanocoumarin, *Liriomyza trifolii*, 519
phototoxicity, 519

G
Galleria mellonella, *Apis mellifera*, 760, 766
microbial control, 760, 766
gamma irradiation, *Ceratitis capitata*, 1449
quarantine, 1449
gamma radiation, *Citrus*, 905
Pantomorus cervinus, 905
generation time, pesticide resistance, 1184
resistance risk assessment, 1184
genetic control, backcross sterility, 405
tobacco budworm, 405
genetic improvement, resistance, 89
Trioxys Pallidus, 89
genetics, *Haematobia irritans*, 1718
lindane resistance, 1745
permethrin, 1718
Tribolium castaneum, 1745
genitalia, *Blatta orientalis*, 1415
hydroprene, 1415
geostatistics, sample patterns, 1888
spatial analysis, 1888
Gleditsia triacanthos, phenology, 869
sampling, 869
glucosinolates, *Lygus* spp., 2258
oilseed rape, 2258
Glycine max, *Anticarsia gemmatalis*, 2107
combined injuries, 599
host plant resistance, 2107
Plathypena scabra, 599
Goniozus legneri, Bethylidae, 733
insecticidal toxins, 733
goodness-of-fit, dose mortality, 1649
extravariation, 1649
Gossypium, spider mites, 1604
Tetranychus, 1604
gossyplure, *Pectinophora gossypiella*, 1321
pheromones, 1321
grain sorghum, plant resistance, 2476
Schizphis graminum, 2476
granaries, chlorpyrifos-methyl, 1128
oats, 1128
grape phylloxera, biotypes, 489
Daktulosphaira vitifoliae, 489, 494
indoleacetic acid, 494

grasses, foraging, 260
Lotus corniculatus, 260
grasshoppers, feeding, 2421
insecticides and temperature, 366
kochia, 2421
pyrethroids, 366
growth, *Blattella germanica*, 2295
insect growth regulators, 2295
gypsy moth maternal effects, *Bacillus thuringiensis*, 2211
microbial pesticide resistance, 2211
gypsy moths, aerial deposition, 173
Bacillus thuringiensis, 173

H
Haematobia irritans, cattle, 1430
ear tags, 1715
genetics, 1718
insecticide metabolism, 662
insecticide resistance, 690
integrated pest management, 1430
permethrin, 1718
pyrethroid resistance, 662, 1715
resistance management, 690
heat, *Dacus dorsalis*, 160
fruit flies, 1959
papayas, 160
quarantine treatments, 1959
heat accumulation, *Malacosoma californicum*, 1492
seasonal history, 1492
heat treatment, *Anastrepha suspensa*, 2331
quarantine security, 2331
Helianthus annuus, *Homoeosoma electellum*, 2219
phenolic acids, 2219
Helicoverpa armigera, pyrethroids, 705, 1682
resistance, 705, 1682
Helicoverpa spp., cotton, 1827
diapause, 1827
Heliothis, host selection, 398
mass rearing, 398
pyrethroids, 1207
resistance, 1207
Heliothis virescens, cell culture, 737
continuous measure, 1222
cotton, 737
insecticide resistance, 1222
parasitoids, 1777
plant allelochemical, 1777
pyrethroids, 335
resistance, 335
Heliothis zea, antibiosis, 614
cotton, 2115
population reduction, 614
toxicity, 2115
herbivory, fecundity, 377
necrophagy, 377
Heterorhabditis, antidesiccants, 1795
Steinernema, 1795
hexythiazox, avermectin, 1752
fenbutatin oxide, 1752
Hippodamia variegata, *Coccinella septempunctata*, 1292
Propylea quatuordecimpunctata, 1292
Homadaula anisocentra, host plant resistance, 533
urban entomology, 533
Homoeosoma electellum, *Helianthus annuus*, 2219
phenolic acids, 2219
Homoptera, sampling, 937
trap, 937
honey bees, *Apis mellifera*, 81
feral honey bee colonies, 81
host adaptation, host suitability, 2393
Thyridopteryx ephemeraeformis, 2393
host plant, *Ceratitis*, 1863
infestation rate, 1863
host plant resistance, *Anticarsia gemmatalis*, 2107
Diatraea saccharalis, 221
Glycine max, 2107
Homadaula anisocentra, 533
Phyllophaga congrua, 1558
Schizaphis graminum, 234
small grains, 1558
Sorghum bicolor, 234
stored grain, 1122
stored-product insects, 1122
sugarcane, 221
urban entomology, 533
host plants, cereal aphids, 846
Diuraphis noxia, 846
host selection, *Heliothis*, 398
mass rearing, 398
host suitability, host adaptation, 2393
Thyridopteryx ephemeraeformis, 2393
host–insect interaction, *Epiphyas postvittana*, 2124
resistance, 2124
house fly, formamidine, 2181
synergism of pyrethroids, 2181
humidity, controlled atmosphere, 472
Cydia pomonella, 472
Humulus lupulus, *Phorodon humuli*, 1388
sampling, 1388
hydramethylnon, fenoxycarb, 74
Pheidole megacephala, 74
hydroprene, *Blatta orientalis*, 1415
genitalia, 1415
hydroxamic acids, *Diadegma terebrans*, 356
Ostrinia nubilalis, 356

I
immigration, *Empoasca fabae*, 1858
Medicago sativa, 1858
indoleacetic acid, *Daktulosphaira vitifoliae*, 494
grape phylloxera, 494
infestation rates, blossom end defects, 1883
Ceratitis, 1863
Dacus, 476, 1883
host plant, 1863
ripeness index, 476
insect control, aesthetic injury level, 2044
Anisota senatoria, 2044
birdsfoot trefoil, 2086
Nematoda, 107
Otiorhynchus sulcatus, 107
plant bugs, 2086
insect exclusion, Lepidoptera, 948
row covers, 948
insect growth regulators, *Blattella germanica*, 2295
cyromazine, 2398
Delia antiqua, 2398
growth, 2295
insect pheromone, disparlure, 910
Lymantria dispar, 910
insect traps, aggregation pheromones, 273
Oryzaephilus surinamensis, 273
insect–plant interactions, *Anasa tristis*, 943
Cucurbitaceae, 943
Eriosoma lanigerum, 1526
integrated pest management, 1526
insect-resistant packaging, *Bracon hebetor*, 1110
Cadra cautella, 1110
insecticide resistance, aphids, 41
filberts, 41
insecticidal baits, *Blattella germanica*, 153
Ebeling choice box, 153
insecticidal toxins, Bethylidae, 733
Goniozus legneri, 733
insecticide bark bands, elm, 1495
Xanthogaleruca luteola, 1495
insecticide metabolism,

Haematobia irritans, 662
Pseudoplusia includens, 672
pyrethroid resistance, 662
resistance, 672
insecticide resistance, *Bacillus thuringiensis*, 1671
Bemisia tabaci, 1665
Colorado potato beetles, 315
continuous measure, 1222
crop rotation, 315
Culex quinquefasciatus, 1704
dieldrin, 1733
Drosophila melanogaster, 1733
esterase activity, 1665
esterases, 1704
Haematobia irritans, 690
Heliothis virescens, 1222
Liriomyza trifolii, 18
microbial pesticide, 1671
permethrin, 1211
Plutella, 1211
resistance management, 690
stored corn, 325
synergism, 18
Typhaea stercorea, 325
insecticide rotation, resistance management, 306
Scirtothrips citri, 306
insecticide selectivity, abamectin, 1783
integrated pest management, 1783
insecticide susceptibility, *Leptinotarsa decemlineata*, 666
insecticides, *Aonidiella aurantii*, 189
behavior, 217
behavioral resistance, 1216
Blattella germanica, 142, 2290
corn, 1114
corn rootworms, 226
dispersal, 142
Lepidoptera, 1002
nematicides, 1002
phenology, 189
pyrethroids, 217, 1216
residual, 226
resistance, 2147
Reticulitermes hesperus, 1395
Schizaphis graminum, 2147
stored-product insects, 1114
toxicity, 2290
tunneling, 1395
insecticides and temperature, grasshoppers, 366
pyrethroids, 366
integrated mite management, pheromone trapping, 985
Phyllonorycter blancardella, 985
integrated pest management, abamectin, 1783
action threshold, 1381
anthurium, 1531
aphids, 1381
apple pests, 2405
cattle, 1430
Diatraea saccharalis, 1563
economic threshold, 641
Eriosoma lanigerum, 1526
floriculture, 1531
Frankliniella occidentalis, 1519
Haematobia irritans, 1430
insect–plant interactions, 1526
insecticide selectivity, 1783
Liriomyza spp., 2153
pesticide resistance, 2153
plant resistance, 1563
Rhagoletis pomonella, 2405
soybeans, 641
Thrips palmi, 1519
Ipomoea batatas, controlled atmosphere storage, 461
Cylas formicarius elegantulus, 461, 1901
sampling, 1901
Iridomyrmex humilis, barrier treatments, 1402
repellency, 1402
irradiation, disinfestation, 1455
Tetranychus urticae, 1455

J

juvenile hormone, *Cacopsylla pyricola*, 293
overwintering, 293
juvenile hormone mimic, reproductive behavior, 2263
mimic, *Trichoplusia ni*, 2263

K

Keiferia lycopersicella, chemical control, 212
tomatoes, 212
Kentucky bluegrass, Cryptostigmata, 205
Poa pratensis, 205
soil ecology, 2362
turfgrass, 2362
kiwifruit, *Dacus tryoni*, 2317
quarantine treatments, 2317
knockdown, *Apis mellifera*, 1925
wasp and hornet sprays, 1925
knockdown resistance, *Musca domestica*, 686
pyrethroid, 686
kochia, feeding, 2421
grasshoppers, 2421

L

Labidura riparia, cultural control, 1034
Elasmopalpus lignosellus, 1034
laboratory bioassays, *Agrotis ipsilon*, 2073
pyrethroids, 2073
Lagenidium giganteum, oomycetes, 374
safety tests, 374
leaf abscission, plant growth, 995
Unaspis euonymi, 995
leafminers, life history, 117
temperature, 117
learning, biological control, 126
Exeristes roborator, 126
lectins, *Diabrotica undecimpunctata howardi*, 2480
Ostrinia nubilalis, 2480
Lepidoptera, cauliflower, 1499
Cochylis hospes, 135
defoliation, 1499
insect exclusion, 948
insecticides, 1002
nematicides, 1002
rearing, 1535
rearing equipment, 1535
row covers, 948
sunflowers, 135
Leptinotarsa decemlineata, *Coleomegilla maculata*, 1306
economic injury level, 2058
insecticide susceptibility, 666
Phalangium opilio, 772
potatoes, 1982, 2058
predation, 1306
resistance, 1229
sampling precision, 428
straw mulch, 1982
life history, leafminers, 117
temperature, 117
life tables, *Myzus nicotianae*, 1080
pest control timing, 1143
population growth, 1080
simulation, 1143
limes, economic injury level, 2008
Polyphagotarsonemus latus, 2008
linalool, *Ceratitis capitata*, 2235
pyrazine, 2235
lindane resistance, genetics, 1745
Tribolium castaneum, 1745
Liriomyza trifolii, biological control, 2380
chrysanthemums, 1246
entomogenous nematodes, 2380
furanocoumarin, 519
insecticide resistance, 18
neem seed extract, 1246
phototoxicity, 519
synergism, 18
Liriomyza spp., integrated pest management, 2153
pesticide resistance, 2153
longevity, fecundity, 1333
Thyanta pallidovirens, 1333
Lotus corniculatus, foraging, 260
grasses, 260

low temperature, *Empoasca fabae*, 1541
mortality response, 1541
lure, attractant, 1350
Ceratitis capitata, 1350
Diabrotica barberi, 1316
disubstituted benzenes, 1316
Lygus hesperus, esterases, 725
resistance, 725
Lygus lineolaris, feeding duration, 2443
population density, 2443
Lygus spp., bioassay methods, 2200
glucosinolates, 2258
oilseed rape, 2258
thuringiensin, 2200
Lymantria dispar, avermectin, 710, 917
biological control, 2229
disparlure, 910, 1977
insect pheromone, 910
mating disruption, 1972
MK-243, 917
pheromones, 1972
predators, 2229
rubidium, 2343
sex pheromones, 1977
sublethal, 710
tree injection, 2343

M

Macrodactylus subspinosus, attractancy, 111
color preference, 111
Magicicada septendecim, apples, 2401
control, 2401
maize, root damage, 2414
western corn rootworm, 2414
Malacosoma californicum, heat accumulation, 1492
seasonal history, 1492
malathion, chlorpyrifos methyl, 1636
rice, 1636
Mallophaga, Anoplura, 1435
sampling, 1435
Malus domestica, *Panonychus ulmi*, 2026
yield, 2026
Mammalia, *Bacillus sphaericus*, 347
Bacillus thuringiensis ssp israelensis, 347
mange, pruritus, 1439
swine, 1439
Mangifera indica, *Anastrepha suspensa*, 2320
quarantine security, 2320
marine areas, *Stomoxys calcitrans*, 887
survival, 887
mark-release-capture, *Diabrotica virgifera virgifera*, 1360
tethered-flight assay, 1360
mass rearing, *Heliothis*, 398
host selection, 398
mating, oviposition, 1837
Pectinophora gossypiella, 1837
mating disruption, *Lymantria dispar*, 1972
pheromones, 1972
maximum pest limit, quarantine security, 13
Tephritidae, 13
Mayetiola destructor, biological control, 2269
cultural control, 1025
degree-days, 1015
parasitoids, 2269
phenology, 1015
plant resistance, 1135
Triticum aestivum, 1025
wheat, 1135
Medicago sativa, *Empoasca fabae*, 1858
immigration, 1858
Melanoplus sanguinipes, biotic potential, 1338
food preference, 1338
metabolic heat, critical radius, 896
thermal conductivity, 896
metabolism, carbendazim, 742
propargite, 63
Rhizoglyphus echinopus, 63
Rhizoglyphus echinopus, 742
metamorphosis, azadirachtin, 2168
Tephritidae, 2168
methomyl, *Phyllonorycter blancardella*, 320
resistance, 320
methoprene, fruit flies, 901
papaya, 901
methyl parathion, chlordimeform, 1549
cotton phytotoxicity, 1549
microbial control, *Apis mellifera*, 760, 766
Beauveria bassiana, 1276
Chalcodermus bimaculatus, 1276
Galleria mellonella, 760, 766
microbial pesticide, *Bacillus thuringiensis*, 1671
insecticide resistance, 1671
microbial pesticide resistance, *Bacillus thuringiensis*, 2211
gypsy moth maternal effects, 2211
microbial safety test, *Apis mellifera*, 755
nontarget organism, 755
mirid pests, biological control, 101
parasites, 101
mites, pears, 2032
sampling, 2032
MK-243, abamectin, 917
Lymantria dispar, 917
models, multiple alleles, 1170
population dynamics, 2469
resistance management, 1170
temperature, 2469
Monellia caryella, pecans, 1801
triphenyltin hydroxide, 1801
monitoring, *Choristoneura rosaceana*, 434
Cydia pomonella, 434
monoterpenes, *Peridroma saucia*, 1761
toxicity, 1761
mortality, carbaryl consumption, 2164
wild rodents, 2164
mortality response, *Empoasca fabae*, 1541
low temperature, 1541
mulch, *Anasa tristis*, 1988
vegetable production systems, 1988
multiple alleles, model, 1170
resistance management, 1170
Musca domestica, abamectin, 1935
bioassays, 1935
control, 788
knockdown resistance, 686
Muscovy ducks, 788
organophosphate resistance, 48
parasites, 799
pyrethroid, 686
resistance genetics, 48
Stomoxys calcitrans, 799
Muscidae, bovine dung, 1446
nematodes, 1446
Muscovy ducks, control, 788
Musca domestica, 788
museum pest control, Anobiidae, 879
Dermestidae, 879
mutagenicity, *Drosophila melanogaster*, 721
permethrin, 721
Myzus nicotianae, esterase, 2130
life tables, 1080
population growth, 1080
resistance, 2130
Myzus persicae, oil, 2352
sequential sampling, 1365
Taylor's power law, 1365
turnip mosaic virus, 2352

N

Nabis spp., predators, 823
seasonal occurrence, 823
necrophagy, fecundity, 377
herbivory, 377

neem seed extract, chrysanthemums, 1246
Liriomyza trifolii, 1246
nematicides, insecticides, 1002
Lepidoptera, 1002
Nematoda, biological control, 1286
insect control, 107
Otiorhynchus sulcatus, 107
Steinernema feltiae, 1286
nematodes, biological control, 1818
bovine dung, 1446
Cylas formicarius elegantulus, 1818
Muscidae, 1446
Neobellieria bullata, 20-OH ecdysone, 2175
bipyridylium herbicides, 2175
Ninhydrin, amino acid, 1633
spray deposition, 1633
nitidulid beetles, date palms, 2357
Ectomyelois ceratoniae, 2357
Nomuraea rileyi, fungicide, 2193
soybean defoliators, 2193
nonpreference, *Tetranychus urticae*, 500
tomatoes, 500
nontarget effects, persistence, 1768
S-31183, 1768
nontarget organism, *Apis mellifera*, 755
microbial safety test, 755

O

oats, chlorpyrifos-methyl, 1128
granaries, 1128
Oebalus pugnax, residual activity, 591
rice, 591
offspring-parent regression, sibling analysis, 647
threshold trait analysis, 647
oil, *Myzus persicae*, 2352
turnip mosaic virus, 2352
oilseed rape, glucosinolates, 2258
Lygus spp., 2258
Oligonychus pratensis, chemical control, 1621
chlorpyrifos, 1570
corn, 1069, 1621
economic injury level, 1069
parathion, 1570
oomycetes, *Lagenidium giganteum*, 374
safety tests, 374
Operophtera brumata, *Bacillus thuringiensis*, 920
Panonychus ulmi, 920
organophosphate resistance, *Culex* spp., 2187
esterases, 2187
Musca domestica, 48
resistance genetics, 48
organophosphates, resistance monitoring, 1194
sex pheromone traps, 1194
Orgyia pseudotsugata, disruption, 1487
pheromones, 1487
Orthoptera, *Apis mellifera*, 784
Poxviridae, 784
Oryzaephilus surinamensis, aggregation pheromones, 273
insect traps, 273
Ostrinia nubilalis, *Bacillus thuringiensis*, 1813
bioassay, 1049
Ostrinia nubilalis, chemigation, 1049
conservation tillage, 2455
corn, 1595
Diabrotica undecimpunctata howardi, 2480
Diadegma terebrans, 356
dispersal, 831
economic injury levels, 1595
feeding stimulant, 2207
formulation, 1813
hydroxamic acids, 356
lectins, 2480
scouting, 831
spray distribution, 2016
starch encapsulation, 2207
vegetable crop, 2016
Otiorhynchus [illegible]stici, alfalfa, 1612
economic injury levels, 1612
Otiorhynchus sulcatus, insect control, 107
Nematoda, 107
overwintering, *Cacopsylla pyricola*, 293
juvenile hormone, 293
oviposition, *Ancylis comptana*, 1514
citrus, 860
codling moths, 131
Cydia pomonella, 131
Diatraea grandiosella, 1575
mating, 1837
Pantomorus cervinus, 860
Pectinophora gossypiella, 1837
Popillia japonica, 2036
Scarabaeidae, 1355
soil pH, 2036
Spodoptera frugiperda, 1575
strawberry, 1514
sugarcane, 1355

P

packaged bees, *Apis mellifera*, 1271
pheromones, 1271
Panonychus citri, *Citrus*, 976
yield, 976
Panonychus ulmi, apples, 552
Bacillus thuringiensis, 920
binomial sampling, 420
economic injury level, 552
Panonychus, acaricides, 1711
Tetranychus, 1711
Panonychus ulmi, European red mite, 547
Malus domestica, 2026
Operophtera brumata, 920
residual bioassay, 961
resistant germ plasma, 180
stress, 547
Tetranychus urticae, 420, 961
Venturia inaequalis, 180
yield, 2026
Pantomorus cervinus, citrus, 860, 905
gamma radiation, 905
oviposition, 860
Papaipema nebris, seedling corn, 1582
yield, 1582
papayas, *Dacus dorsalis*, 160, 1944
Dacus spp., 2327
fruit flies, 901
heat, 160
methoprene, 901
quarantine treatment, 2327
temperature stress, 1944
Parapediasia teterrella, sampling, 856
sex pheromones, 856
parasites, biological control, 101, 1096
mirid pests, 101
Musca domestica, 799
Stomoxys calcitrans, 799
stored-product insects, 1096
parasitism, sorghum, 84
stem borers, 84
parasitoids, biological control, 2269
Heliothis virescens, 1777
Mayetiola destructor, 2269
plant allelochemical, 1777
parathion, chlorpyrifos, 1570
Oligonychus pratensis, 1570
peach injury, flower thrips, 511
sampling techniques, 511
pearl millet, plant diseases, 2102
varietal resistance, 2102
pears, mites, 2032
sampling, 2032
pecan aphids, pesticides, 1806
predators, 1806
pecans, *Monellia caryella*, 1801
triphenyltin hydroxide, 1801
Pectinophora gossypiella, chemical termination treatment, 1074
cotton, 1074

gossyplure, 1321
mating, 1837
oviposition, 1837
pheromones, 1321
Peridroma saucia, monoterpenes, 1761
toxicity, 1761
Perkinsiella saccharicida, sequential sampling, 2284
Tytthus, 2284
permethrin, *Drosophila melanogaster*, 721
genetics, 1718
Haematobia irritans, 1718
insecticide resistance, 1211
mutagenicity, 721
Plutella, 1211
persistence, nontarget effects, 1768
S-31183, 1768
pest control, chemical adsorption, 55
field burning, 55
pest control timing, life table, 1143
simulation, 1143
pest management, *Anthonomus eugenii*, 2003
Campylomma verbasci, 1506
Capsicum annuum, 2003
Delia coarctata, 2065
expert system, 2065
pheromones, 1506
sampling, 1656
Taylor coefficients, 1656
pest resurgence, dispersion, 1847
Tetranychus urticae, 1847
pesticide resistance, generation time, 1184
integrated pest management, 2153
Liriomyza spp., 2153
resistance risk assessment, 1184
pesticide spray distribution, water-soluble dyes, 2411
pesticides, Arachnida, 8
bioassays, 8
pecan aphids, 1806
predators, 1806
Phalangium opilio, *Leptinotarsa decemlineata*, 772
Pheidole megacephala, fenoxycarb, 74
hydramethylnon, 74
phenolic acids, *Helianthus annuus*, 2219
Homoeosoma electellum, 2219
phenolics, strawberries, 564
strawberries, 557
Tetranychus urticae, 557, 564
phenology, *Aonidiella aurantii*, 189
degree-days, 1915
Gleditsia triacanthos, 869
insecticides, 189
Mayetiola destructor, 1015
sampling, 869
pheromones, *Anthonomus grandis grandis*, 610
Apis mellifera, 1271
Campylomma verbasci, 1506
Coleoptera, 268
disruption, 1487
gossyplure, 1321
Lymantria dispar, 1972
mating disruption, 1972
Orgyia pseudotsugata, 1487
packaged bees, 1271
Pectinophora gossypiella, 1321
pest management, 1506
stored grain, 268
pheromone trapping, integrated mite management, 985
Phyllonorycter blancardella, 985
pheromone traps, fruit orchards, 197
trap efficiency, 197
Phorodon humuli, *Humulus lupulus*, 1388
sampling, 1388
phosphine, distribution, 1459
fumigant, 1459
resistance, 1738
stored products, 1738
photosynthesis, resistance mechanisms, 2111
Sipha flava, 2111
phototoxicity, furanocoumarin, 519
Liriomyza trifolii, 519
Phyllonorycter blancardella, integrated mite management, 985
methomyl, 320
pheromone trapping, 985
resistance, 320
resistance management, 678
Phyllophaga congrua, host plant resistance, 1558
small grains, 1558
pistachios, *Calocoris norvegicus*, 1841
weeds, 1841
plant allelochemical, *Heliothis virescens*, 1777
parasitoids, 1777
plant bugs, birdsfoot trefoil, 580, 2086
insect control, 2086
plant damage, 580
plant damage, birdsfoot trefoil, 580
plant bugs, 580
plant diseases, pearl millet, 2102
varietal resistance, 2102
plant growth, leaf abscission, 995
Unaspis euonymi, 995
plant resistance, antibiosis, 1053
Blissus leucopterus hirtus, 1640
Diatraea grandiosella, 1578
Diatraea saccharalis, 866, 1563
endophyte, 1640
grain sorghum, 2476
integrated pest management, 1563
Mayetiola destructor, 1135
pubescence, 866
Schizaphis graminum, 2476
Sipha flava, 1053
Spodoptera frugiperda, 1578
wheat, 1135
Plathypena scabra, combined injuries, 599
Glycine max, 599
Platynota idaeusalis, resistance monitoring, 329
sex pheromone traps, 329
Plutella, insecticide resistance, 1211
permethrin, 1211
Plutella xylostella, cabbage, 1201
resistance, 1201
Poa pratensis, Cryptostigmata, 205
Kentucky bluegrass, 205
Polyphagotarsonemus latus, economic injury level, 2008
limes, 2008
Popillia japonica, oviposition, 2036
soil pH, 2036
population density, feeding duration, 2443
Lygus lineolaris, 2443
population dynamics, *Apis mellifera*, 1791
digitizer, 1791
models, 2469
temperature, 2469
population effects, *Blattella germanica*, 444
fenoxycarb, 444
population estimates, *Apis mellifera*, 748
colony strength, 748
population growth, life tables, 1080
Myzus nicotianae, 1080
population management, eradication, 1344
quarantine, 1344
population monitoring, almonds, 570
Amyelois transitella, 570

population reduction, antibiosis, 614
Heliothis zea, 614
postharvest quality, *Cydia pomonella*, 2335
Prunus spp., 2335
potatoes, economic injury level, 2058
Leptinotarsa decemlineata, 1982, 2058
straw mulch, 1982
poultry, flies, 451
predators, 451
Poxviridae, *Apis mellifera*, 784
Orthoptera, 784
predation, *Anthonomus grandis*, 1
Coleomegilla maculata, 1306
Leptinotarsa decemlineata, 1306
Solenopsis invicta, 1
predators, biological control, 2229
flies, 451
Lymantria dispar, 2229
Nabis spp., 823
pecan aphids, 1806
pesticides, 1806
poultry, 451
seasonal occurrence, 823
probe traps, Coleoptera, 1102
distribution, 1102
propargite, metabolism, 63
Rhizoglyphus echinopus, 63
Tetranychus pacificus, 655
Tetranychus urticae, 655
propoxur, *Blattella germanica*, 1912
chlorpyrifos, 1912
Propylea quatuordecimpunctata, *Coccinella septempunctata*, 1292
Hippodamia variegata, 1292
protein, *Ceratitis capitata*, 1954
diet, 1954
Prunus, apples, 466
Cydia pomonella, 466
Prunus spp., *Cydia pomonella*, 2335
postharvest quality, 2335
pruritus, mange, 1439
swine, 1439
Pseudaletia unipuncta, *Agrotis ipsilon*, 596
sampling, 596
Pseudoplusia includens, field control, 27
insecticide metabolism, 672
pyrethroids, 35
resistance, 35, 672
soybean cyst nematode, 247
soybeans, 27
stem canker fungus, 247
Pseudotsuga menziesii, cone and seed insects, 1485
western conifer seed bug, 1485
Psylla pyricola, pyrethroids, 1177
resistance management, 1177
pteridine eye pigments, age distribution, 1422
Cochliomyia hominivorax, 1422
pubescence, *Diatraea saccharalis*, 866
plant resistance, 866
Pyemotes tritici, reproduction, 384
stored products, 384
pyrazine, *Ceratitis capitata*, 2235
linalool, 2235
pyrethroid resistance, ear tags, 1715
Haematobia irritans, 662, 1715
insecticide metabolism, 662
Scirtothrips citri, 1723
synergism, 1723
pyrethroids, *Agrotis ipsilon*, 2073
behavior, 217, 1510
behavioral resistance, 1216
Cacopsylla pyricola, 2159
carbamate, 2136
grasshoppers, 366
Helicoverpa armigera, 705, 1682
Heliothis, 1207
Heliothis virescens, 335
insecticides, 217, 1216
insecticides and temperature, 366
knockdown resistance, 686
laboratory bioassays, 2073
Musca domestica, 686
Pseudoplusia includens , 35
Psylla pyricola, 1177
resistance, 35, 335, 705, 1207, 1682, 2159
resistance management, 1177
Spodoptera exigua, 2136
temperature coefficient, 342
Trichoplusia ni, 342, 1510
Pyrrhalta, biological control, 2196
diapause, 2196

Q

Quadraspidiotus perniciosus, commodity, 892
radiation, 892
quarantine, *Ceratitis capitata*, 1449
detector dog, 1932
eradication, 1344
gamma irradiation, 1449
population management, 1344
screwworms, 1932
quarantine security, *Anastrepha suspensa*, 1468, 1471, 1475, 2320, 2331, 2340
commodity treatment, 1471, 1475, 2340
fruit quality, 1468
heat treatment, 2331
Mangifera indica, 2320
maximum pest limit, 13
Tephritidae, 13
quarantine treatments, *Anastrepha*, 1940
Anastrepha suspensa, 2324
Ceratitis, 1940
Dacus spp., 2327
Dacus tryoni, 2317
fruit flies, 1959, 1965
heat, 1959
kiwifruit, 2317
papayas, 2327
shrink wrapping, 2324
vapor heat, 1965

R

radiation, commodity, 892
Quadraspidiotus perniciosus, 892
rapeseed, *Brassica napus* L., 2246
rearing, Lepidoptera, 1535
rearing equipment, 1535
rearing equipment, Lepidoptera, 1535
rearing, 1535
repellency, barrier treatments, 1402
Coptotermes formosanus, 875
Ctenocephalides felis felis, 2306
Felis catus, 2306
Iridomyrmex humilis, 1402
termiticides, 875
repellents, feeding deterrents, 629
Rhyzopertha dominica, 629
reproduction, *Blattella germanica*, 441
boll weevils, 392
chlorpyrifos, 441
Pyemotes tritici, 384
Sphaeralcea spp., 392
stored products, 384
reproductive behaviors, juvenile hormone mimic, 2263
Trichoplusia ni, 2263
residual, corn rootworms, 226
insecticides, 226
residual activity, *Oebalus pugnax*, 591
rice, 591
residual bioassay, *Panonychus ulmi*, 961
Tetranychus urticae, 961
residual insecticides, *Blattella germanica*, 1907
crack and crevice, 1907
resistance, *Apis mellifera*, 715
cabbage, 1201
Cacopsylla pyricola, 2159

Clavigralla tomentosicollis, 300
cockroaches, 1698
Cynodon dactylon, 585
cyromazine, 1689
detoxification, 715
diagnostic dose, 1151, 1160
diflubenzuron, 1689
Diuraphis noxia, 1091
Epiphyas postvittana, 2124
esterases, 725, 2130
genetic improvement, 89
Helicoverpa armigera, 705, 1682
Heliothis, 1207
Heliothis virescens, 335
host–insect interaction, 2124
insecticide metabolism, 672
insecticides, 2147
Leptinotarsa decemlineata, 1229
Lygus hesperus, 725
methomyl, 320
Myzus nicotianae, 2130
phosphine, 1738
Phyllonorycter blancardella, 320
Plutella xylostella, 1201
Pseudoplusia includens , 35, 672
pyrethroids, 35, 335, 705, 1207, 1682, 2159
Schizaphis graminum, 2147
spider mites, 1236
Spodoptera frugiperda, 585
Spodoptera littoralis, 59
statistical power, 1160
statistical test power, 1151
stored products, 1738
synergism, 1236
synergist, 1698
teflubenzuron, 59
Trioxys Pallidus, 89
wheat, 1091
wild cowpea, 300
resistance genetics, *Musca domestica*, 48
organophosphate resistance, 48
resistance management, *Haematobia irritans*, 690
insecticide resistance, 690
insecticide rotation, 306
model, 1170
multiple alleles, 1170
Phyllonorycter blancardella, 678
Psylla pyricola, 1177
pyrethroids, 1177
Scirtothrips citri, 306
resistance mechanisms, photosynthesis, 2111
Sipha flava, 2111
resistance monitoring, organophosphates, 1194
Platynota idaeusalis, 329
sex pheromone traps, 329, 1194
resistance risk assessment, generation time, 1184
pesticide resistance, 1184
resistant germ plasma, *Panonychus ulmi*, 180
Venturia inaequalis, 180
Reticulitermes flavipes, *Coptotermes formosanus*, 1918
tunneling response, 1918
Reticulitermes hesperus, insecticides, 1395
tunneling, 1395
Rhabditida, biological control, 97
entomogenous nematodes, 97
Rhagoletis mendax, risk prediction, 526
Vaccinium spp., 526
Rhagoletis pomonella, apple pests, 2405
apple volatiles, 539
integrated pest management, 2405
trapping, 539
Rhizoglyphus echinopus, metabolism, 63
propargite, 63
Rhizoglyphus echinopus, carbendazim, 742
metabolism, 742
Rhopalosiphum maidis, corn, 1626
DIMBOA, 1626
Rhyzopertha dominica, feeding deterrents, 629
fumigants, 1677
repellents, 629
sampling, 625
stored products, 625
Tribolium castaneum, 1677
rice, chlorpyrifos methyl, 1636
malathion, 1636
Oebalus pugnax, 591
residual activity, 591
ripeness index, *Dacus*, 476
infestation rates, 476
risk prediction, *Rhagoletis mendax*, 526
Vaccinium spp., 526
root damage, maize, 2414
western corn rootworm, 2414
row covers, insect exclusion, 948
Lepidoptera, 948
rubidium, *Lymantria dispar*, 2343
tree injection, 2343
Rutelinae, attractants, 1298
Cetoniinae, 1298

S

S-31183, nontarget effects, 1768
persistence, 1768
safety tests, *Lagenidium giganteum*, 374
oomycetes, 374
sample patterns, geostatistics, 1888
spatial analysis, 1888
sampling, *Agrotis ipsilon*, 596
Anarsia lineatella, 505
Anoplura, 1435
corn rootworms, 574
Cylas formicarius elegantulus, 1901
Dacus cucurbitae, 2274
Dacus dorsalis, 2274
Diabrotica, 574
Diatraea grandiosella, 1370
distribution, 505
Gleditsia triacanthos, 869
Homoptera, 937
Humulus lupulus, 1388
Ipomoea batatas, 1901
Mallophaga, 1435
mites, 2032
Parapediasia teterrella, 856
pears, 2032
pest management, 1656
phenology, 869
Phorodon humuli, 1388
Pseudaletia unipuncta, 596
Rhyzopertha dominica, 625
sex pheromones, 856
stored products, 625
Taylor coefficients, 1656
Taylor's power law, 1370
Tetranychus urticae, 1376
trap, 937
Typhlodromus occidentalis, 1376
sampling precision, *Leptinotarsa decemlineata*, 428
sampling procedures, cotton, 1064
Frankliniella spp., 1064
sampling techniques, flower thrips, 511
peach injury, 511
Scarabaeidae, oviposition, 1355
sugarcane, 1355
Schizaphis graminum, feeding behavior, 241
host plant resistance, 234
insecticides, 2147
resistance, 2147
Sorghum bicolor, 234, 241
grain sorghum, 2476
plant resistance, 2476
Scirtothrips citri, cross-resistance, 698
fluvalinate resistance, 698
insecticide rotation, 306
pyrethroid resistance, 1723

resistance management, 306
synergism, 1723
scouting, dispersal, 831
Ostrinia nubilalis, 831
screwworms, detector dog, 1932
quarantine, 1932
seasonal history, heat accumulation, 1492
Malacosoma californicum, 1492
seasonal occurrence, *Nabis* spp., 823
predators, 823
seed orchards, *Conophthorus coniperda*, 2349
white pine, 2349
seedling corn, *Papaipema nebris*, 1582
yield, 1582
semiochemicals, attraction, 2040
Toxotrypana curvicauda, 2040
sequential sampling, *Myzus persicae*, 1365
Perkinsiella saccharicida, 2284
Taylor's power law, 1365
Tytthus, 2284
settling, acceptance, 814
Empoasca fabae, 814
sex pheromone traps, organophosphates, 1194
Platynota idaeusalis, 329
resistance monitoring, 329, 1194
sex pheromones, disparlure, 1977
Lymantria dispar, 1977
Parapediasia teterrella, 856
sampling, 856
sex ratio, *Biosteres tryoni*, 779
Ceratitis capitata, 779
shrink wrapping, *Anastrepha suspensa*, 2324
quarantine treatment, 2324
sibling analysis, offspring-parent regression, 647
threshold trait analysis, 647
simulation, life table, 1143
pest control timing, 1143
Sipha flava, antibiosis, 1053
economic injury level, 1008
photosynthesis, 2111
plant resistance, 1053
resistance mechanisms, 2111
sorghum, 1008
Sitophilus zeamais, *Anisopteromalus calandrae*, 2462
biological control, 2462
small grains, host plant resistance, 1558
Phyllophaga congrua, 1558
soil, benzodioxole, 1261
fruit flies, 1261
soil ecology, Kentucky bluegrass, 2362
turfgrass, 2362
soil pH, oviposition, 2036
Popillia japonica, 2036
Solenopsis invicta, *Anthonomus grandis*, 1
baits, 67
Diptera pupae, 67
predation, 1
sorghum, economic injury level, 1008
parasitism, 84
Sipha flava, 1008
stem borers, 84
Sorghum bicolor, feeding behavior, 241
host plant resistance, 234
Schizaphis graminum, 234, 241
soybean cyst nematode, *Pseudoplusia includens*, 247
stem canker fungus, 247
soybean defoliators, fungicide, 2193
Nomuraea rileyi, 2193
soybeans, economic threshold, 641
field control, 27
integrated pest management, 641
Pseudoplusia includens, 27
spatial analysis, geostatistics, 1888
sample patterns, 1888
sperm, *Ceratitis capitata*, 1949
distribution, 1949
Sphaeralcea spp., boll weevils, 392
reproduction, 392
spider mites, *Gossypium*, 1604
resistance, 1236
synergism, 1236
Tetranychus, 1604
Spodoptera exigua, carbamate, 2136
pyrethroids, 2136
Spodoptera frugiperda, corn, 2452
Cynodon dactylon, 585
Diatraea grandiosella, 1575, 1578
economic injury, 2452
oviposition, 1575
plant resistance, 1578
resistance, 585
Spodoptera littoralis, resistance, 59
teflubenzuron, 59
spray deposition, amino acid, 1633
Ninhydrin, 1633
spray distribution, *Ostrinia nubilalis*, 2016
vegetable crops, 2016
stalkborers, corn, 850
feeding behavior, 850
starch encapsulation, feeding stimulant, 2207
Ostrinia nubilalis, 2207
statistical power, diagnostic dose, 1160
resistance, 1160
statistical test power, diagnostic dose, 1151
resistance, 1151
Steinernema, antidesiccants, 1795
Heterorhabditis, 1795
Steinernema feltiae, biological control, 1286
Nematoda, 1286
stem borers, parasitism, 84
sorghum, 84
stem canker fungus, *Pseudoplusia includens*, 247
soybean cyst nematode, 247
sterility, *Anthonomus grandis grandis*, 1853
egg hatch, 1853
Stomoxys calcitrans, adhesives, 883
traps, 883
marine areas, 887
Musca domestica, 799
parasites, 799
survival, 887
stored corn, insecticide resistance, 325
Typhaea stercorea, 325
stored grain, Coleoptera, 268
host plant resistance, 1122
pheromones, 268
stored-product insects, 1122
stored products, phosphine, 1738
Pyemotes tritici, 384
reproduction, 384
resistance, 1738
Rhyzopertha dominica, 625
sampling, 625
stored-product insects, biological control, 1096
corn, 1114
host plant resistance, 1122
insecticides, 1114
parasites, 1096
stored grain, 1122
straw mulch, *Leptinotarsa decemlineata*, 1982
potatoes, 1982
strawberries, *Ancylis comptana*, 1514
oviposition, 1514
phenolics, 557, 564
Tetranychus urticae, 557, 564
stress, European red mite, 547
Panonychus ulmi, 547
sublethal, avermectin, 710
Lymantria dispar, 710

sugarbeets, degree-days, 2078
Tetranops myopaeformis, 2078
sugarcane, *Diatraea saccharalis*, 221
host plant resistance, 221
oviposition, 1355
Scarabaeidae, 1355
sulfluramid, *Blattella germanica*, 148, 1409
delayed-action toxicity, 148
toxicity, 1409
sunflowers, *Cochylis hospes*, 135
Lepidoptera, 135
survival, *Anthonomus grandis grandis*, 1879
emergence, 1879
marine areas, 887
Stomoxys calcitrans, 887
swathing, alfalfa, 2427
cutting, 2427
swine, mange, 1439
pruritus, 1439
synergism, insecticide resistance, 18
Liriomyza trifolii, 18
pyrethroid resistance, 1723
resistance, 1236
Scirtothrips citri, 1723
spider mites, 1236
synergism of pyrethroids, formamidine, 2181
house fly, 2181
synergist, cockroaches, 1698
resistance, 1698

T

Taylor coefficients, pest management, 1656
sampling, 1656
Taylor's power law, *Diatraea grandiosella*, 1370
Myzus persicae, 1365
sampling, 1370
sequential sampling, 1365
teflubenzuron, resistance, 59
Spodoptera littoralis, 59
temperature, leafminers, 117
life history, 117
models, 2469
population dynamics, 2469
temperature coefficient, pyrethroids, 342
Trichoplusia ni, 342
temperature stress, *Dacus dorsalis*, 1944
papayas, 1944
Tephritidae, azadirachtin, 2168
maximum pest limit, 13
metamorphosis, 2168
quarantine security, 13
termiticides, *Coptotermes formosanus*, 875
repellency, 875
Tetranops myopaeformis, degree-days, 2078
sugarbeets, 2078
tethered-flight assay, *Diabrotica virgifera virgifera*, 1360
mark-release-capture, 1360
Tetranychus, acaricides, 1711
Gossypium, 1604
Panonychus, 1711
spider mites, 1604
Tetranychus pacificus, *Eotetranychus williamettei*, 2252
propargite, 655
Tetranychus urticae, 655
Vitis vinifera, 2252
Tetranychus urticae, *Bacillus thuringiensis*, 792
binomial sampling, 420
disinfestation, 1455
dispersion, 1847
irradiation, 1455
nonpreference, 500
Panonychus ulmi, 420, 961
pest resurgence, 1847
phenolics, 557, 564
propargite, 655
residual bioassay, 961
sampling, 1376
strawberries, 557, 564
Tetranychus pacificus, 655
thuringiensin, 792
tomatoes, 500
Typhlodromus occidentalis, 1376
thatch, *Blissus leucopterus*, 2370, 2375
turfgrass, 2370, 2375
thermal conductivity, critical radius, 896
metabolic heat, 896
threshold trait analysis, offspring-parent regression, 647
sibling analysis, 647
thrips, cucumbers, 2092
damage, 2092
Thrips palmi, *Frankliniella occidentalis*, 1519
integrated pest management, 1519
thuringiensin, *Bacillus thuringiensis*, 792
bioassay methods, 2200
Lygus spp., 2200
Tetranychus urticae, 792
Thyanta pallidovirens, fecundity, 1333
longevity, 1333
Thyridopteryx ephemeraeformis, host adaptation, 2393
host suitability, 2393
tobacco budworm, backcross sterility, 405
genetic control, 405
tomatoes, chemical control, 212
Keiferia lycopersicella, 212
nonpreference, 500
Tetranychus urticae, 500
toxicity, *Blattella germanica*, 1409, 2290
cotton, 2115
Heliothis zea, 2115
insecticides, 2290
monoterpenes, 1761
Peridroma saucia, 1761
sulfluramid, 1409
Toxotrypana curvicauda, attraction, 2040
semiochemicals, 2040
translocation disruption, alfalfa, 807
Empoasca fabae, 807
trap efficiency, fruit orchards, 197
pheromone traps, 197
trapping, *Anastrepha ludens*, 412
apple volatiles, 539
color, 412
Rhagoletis pomonella, 539
traps, adhesives, 883
Homoptera, 937
sampling, 937
Stomoxys calcitrans, 883
tree injection, *Lymantria dispar*, 2343
rubidium, 2343
Tribolium castaneum, fumigants, 1677
genetics, 1745
lindane resistance, 1745
Rhyzopertha dominica, 1677
Trichoplusia ni, behavior, 1510
juvenile hormone mimic, 2263
pyrethroids, 342, 1510
reproductive behaviors, 2263
temperature coefficient, 342
trimedlure, attractant, 819
Ceratitis capitata, 819
Trioxys Pallidus, genetic improvement, 89
resistance, 89
triphenyltin hydroxide, *Monellia caryella*, 1801
pecans, 1801
Triticum aestivum, cultural control, 1025
Diuraphis noxia, 2434
Mayetiola destructor, 1025
vernalization, 2434
tunneling, insecticides, 1395
Reticulitermes hesperus, 1395
tunneling response, *Coptotermes formosanus*, 1918
Reticulitermes flavipes, 1918
turfgrass, *Blissus leucopterus*, 2370
Blissus leucopterus, 2375
Kentucky bluegrass, 2362
soil ecology, 2362
thatch, 2370, 2375

turnip mosaic virus, *Myzus persicae*, 2352
oil, 2352
20-OH ecdysone, bipyridylium, 2175
Neobellieria bullata, 2175
Typhaea stercorea, insecticide resistance, 325
corn, 325
Typhlodromus occidentalis, sampling, 1376
Tetranychus urticae, 1376
Tytthus, Perkinsiella saccharicida, 2284
sequential sampling, 2284

U
ultraviolet, baculovirus, 168
dyes, 168
Unaspis euonymi, leaf abscission, 995
plant growth, 995
urban entomology, *Homadaula anisocentra*, 533
host plant resistance, 533

V
Vaccinium spp., *Rhagoletis mendax*, 526
risk prediction, 526
vapor heat, fruit flies, 1965
quarantine treatments, 1965
vapor toxicity, *Bemisia tabaci*, 1254
buprofezin, 1254
varietal resistance, pearl millet, 2102
plant diseases, 2102
vegetable crops, *Ostrinia nubilalis*, 2016
spray distribution, 2016
vegetable production systems, *Anasa tristis*, 1988
mulch, 1988
Venturia inaequalis, Panonychus ulmi, 180
resistant germ plasma, 180
vernalization, *Diuraphis noxia*, 2434
Triticum aestivum, 2434
videotape recording, *Aphis gossypii*, 955
bifenthrin, 955
Vitis vinifera, Eotetranychus williamettei, 2252
Tetranychus pacificus, 2252

W
Wald's seq. prob. ratio test, *Choristoneura fumiferana*, 1479
decision support system, 1479
wasp and hornet sprays, *Apis mellifera*, 1925
knockdown, 1925
water-soluble dyes, pesticide spray distribution, 2411
weed competition, *Agrotis ipsilon*, 1058
corn seedling damage, 1058
weeds, *Calocoris norvegicus*, 1841
pistachios, 1841
western conifer seed bug, cone and seed insects, 1485
Pseudotsuga menziesii, 1485
western corn rootworm, maize, 2414
root damage, 2414
western X-disease, *Colladonus montanus*, 2279
Fieberiella florii, 2279
wheat, carbon dioxide, 277
Cryptolestes ferrugineus, 277
Diuraphis noxia, 1091
Mayetiola destructor, 1135
plant resistance, 1135
resistance, 1091
white pine, *Conophthorus coniperda*, 2349
seed orchards, 2349
wild cowpea, *Clavigralla tomentosicollis*, 300
resistance, 300
wild rodents, carbaryl consumption, 2164
mortality, 2164

X
Xanthogaleruca luteola, damage, 837
degree-days, 837
elm, 1495
insecticide bark bands, 1495

Y
yield, *Citrus*, 976
Malus domestica, 2026
Panonychus citri, 976
Panonychus ulmi, 2026
Papaipema nebris, 1582
seedling corn, 1582
yield reduction, *Acrosternum hilare*, 842
cotton, 842

Z
Zea mays, Blissus leucopterus leucopterus, 618